



A FREE PDF

from Steven David Lampley
Stand Up Comedian

What is Laughter?

A Totally Serious Academic
Investigation by Someone Who
Has Laughed Before

What is Laughter?

A Totally Serious [tic] Academic Investigation by Someone Who Has Laughed Before

Laughter is one of the most recognizable human behaviors, yet it remains one of the most complex to define. It is not simply a response to humor, but a multi-modal, social, physiological, and neurochemical event. This paper synthesizes current research from neuroscience, psychology, and physiology to answer what laughter is, how it is produced in the brain and body, and what functions it serves.

ONE: Defining Laughter

Laughter is typically defined as a rhythmic, vocal, everted behavior consisting of short, repeated expirations of air accompanied by characteristic facial expressions and vocalizations. Critically, modern research separates laughter from humor. Long viewed as a byproduct of humor, laughter may have evolved not from punchlines but from mammalian play, according to a growing paradigm in neuroscience. In animals, laughter is a form of social bonding based on a primordial reflex to tickling, but human laughter has come a long way from these playful roots.

Researchers now distinguish between spontaneous, or Duchenne, laughter and conversational or social laughter. These different kinds of laughter also spark different connections within the "laughter perception network" in the human brain depending on their context.

TWO: The Neurobiology of Laughter

The expression of laughter depends on two partially independent neuronal pathways. This dual-pathway model is central to current understanding.

a) The Involuntary Pathway: This older, emotional system originates in subcortical structures including the amygdala, hypothalamus, and periaqueductal gray, with motor effectors in the brainstem. Studies using fMRI during conflicting voluntary vocalization versus laughter show that the laughter-triggering

network relies heavily on sensory and deep interoceptive analysis, as well as on motor effectors in the brainstem. This pathway is responsible for the uncontrollable, contagious laughter that arrests speech.

b) The Voluntary Pathway: This newer system involves premotor and motor cortex, the supplementary motor area, and lateral prefrontal cortex. It allows us to produce polite, social laughter on demand and to inhibit laughter when inappropriate.

c) Perception: Hearing laughter engages a distributed network. Tickle-induced laughter provokes extra activity in the secondary auditory cortex, likely reflecting acoustical complexity, whereas joyful and taunting laughter prompts more activity in the medial frontal cortex, a region associated with social and emotional processing. Individual differences in attributing social intention to laughter are also linked to trait-level differences in mentalizing, with attenuated tendency to attribute positive social intention associated with higher autistic traits.

THREE: Psychological and Social Functions

Psychology has moved from viewing laughter as merely an expression of positive emotion to a functional social signal.

1. Social Bonding Theory: The dominant evolutionary account argues laughter functions to facilitate bonding. In primates, grooming builds bonds via the opioid system. For humans, laughter is faster and gives an enhanced opportunity to build large social networks. Research shows relaxed social laughter is associated with feelings of wellbeing and heightened affect, and that social laughter is correlated with an elevated pain threshold, suggesting an endorphin-mediated opiate effect.

This has been confirmed with neuroimaging. Social laughter triggers endogenous opioid release in humans, and laughter imposes less severe timing constraints on social bonding compared with grooming and could thus play a key role in enabling humans to live in exceptionally large social networks. The relationship between opioid receptor density and laughter rate also suggests the opioid system may underlie individual differences in sociability.

Caruana's emotional mirroring hypothesis adds that a function of laughter is to facilitate emotional contagion and to boost affiliation and social bonding. In the long term, mirroring others' laughter shapes one's own socio-emotional habits.

2. Social Function Differentiation: Recent acoustic work with 762 participants rating 400 laughter samples found that laughter acoustics systematically convey three fundamental social tasks: rewarding behavior, establishing affiliative bonds, and negotiating dominance. Joyful, affiliative laughter is distinct from dominant, mocking laughter in both perception and neural response.

3. Developmental Function: Spontaneous laughter and smiles are the very first prosocial gestures that infants direct toward their caregivers. Laughter activates many of the same brain reward circuits as

food and sex, and a 2025 synthesis finds it measurably lowers cortisol and may restructure how the developing brain builds resilience to stress.

FOUR: The Physiology of Laughter

Laughter is a full-body physiological event, not just a vocalization.

Autonomic and Cardiovascular Effects: Vigorous laughter temporarily increases heart rate, benefits the immune system, increases alertness, and exercises skeletal muscles. Following extended laughter, there is a brief period during which blood pressure lowers and heart rate decreases. It can also stimulate circulation and aid muscle relaxation, both of which can help reduce some of the physical symptoms of stress.

Endocrine and Immune Effects: Humor results in a general decrease in stress hormones. Early work by Berk showed humor reduced cortisol and catecholamine concentrations, which are products of the hypothalamus-pituitary-adrenal axis, as well as increased the production of antibodies and endorphins, the body's natural pain killers.

Laughter therapy physiologically reduces the level of stress hormones, increases health-promoting hormones such as endorphins, and strengthens the immune system by increasing the number of T-lymphocytes through activation of natural killer cells. Lab tests showed that the immune systems of "laughers" tend to have increased quantity and vitality of T cells, and that laughter increases Ig A, G, and M. Even a few minutes of genuine laughter can stimulate changes in immune activity that make the body more resistant to disease.

Long-term, laughter may improve immune system function, as negative thoughts manifest into chemical reactions that can affect the body by bringing more stress and decreasing immunity.

Conclusion

What is laughter? Biologically, it is a dual-pathway motor program rooted in ancient play circuitry and modulated by higher cortical control. Psychologically, it is a sophisticated social signal that rewards, affiliates, and negotiates status. Physiologically, it is a brief stressor that produces a longer-lasting relaxation and immuno-enhancing response.

It is not simply the consequence of understanding a joke. It is a mechanism for maintaining social cohesion, regulating stress, and communicating emotional safety without words. As Dunbar and others argue, human societies became too large for physical grooming to maintain bonds; laughter became vocal grooming — efficient, contagious, and neurochemically rewarding.

In that sense, laughter is best understood not as an emotion itself, but as a regulator of emotions, both our own and those around us.

Bibliography

Wildgruber, D., et al. (2013). Different Types of Laughter Modulate Connectivity within Distinct Parts of the Laughter Perception Network. PLOS ONE. Foundational study showing distinct neural networks for tickling vs. joyful/taunting laughter.

Caruana, F. (2021). Cracking the laugh code: laughter through the lens of biology, psychology and neuroscience. Philosophical Transactions of the Royal Society B. Comprehensive theme issue overview of dual-pathway model.

Scott, S. K., et al. (2014). Neural correlates of laughter and humour. Brain. Review of lesion and stimulation studies defining the two partially independent pathways for laughter expression.

Dunbar, R. I. M., et al. (2012). Social laughter is correlated with an elevated pain threshold. Proceedings of the Royal Society B. Proposes endorphin-mediated bonding function.

Manninen, S., et al. (2017). Social Laughter Triggers Endogenous Opioid Release in Humans. Journal of Neuroscience. PET evidence for opioid release during social laughter.

Wood, A., et al. (2017). Towards a social functional account of laughter: Acoustic features convey reward, affiliation, and dominance. Study of 762 participants rating 400 laughter samples.

Berk, L. S., et al. (1989-2001). Neuroendocrine and stress hormone changes during mirthful laughter. Foundational work showing reduction in cortisol and catecholamines and increase in antibodies and endorphins.

Bennett, M. P. & Lengacher, C. (2008). Humor and Laughter May Influence Health IV. Humor and Immune Function. Review of effects on natural killer cell cytotoxicity and immunoglobulins.

Harding, J. (2025). Analysis in Routledge volume on early childhood, synthesizing neuroscience, developmental psychology, and endocrinology on laughter and resilience.

Mayo Clinic (2023). Mayo Mindfulness: Laughter for stress relief. Clinical summary of circulation, muscle relaxation, and long-term immune benefits.

